



Ab Initio Investigation of the Ground State of C_3H^- : Potential Energy Function and Rovibrational Spectrum

M. Hochlaf, P. Rosmus, N.M. Lakin, J.P. Maier

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 37, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

***Ab initio* investigation of the ground state of C_3H^- : Potential energy function and rovibrational spectrum**

M. HOCHLAF, P. ROSMUS,

Theoretical Chemistry Group, Université de Marne–La–Vallée, F–77454 Champs–sur Marne,
France.

N. M. LAKIN, J. P. MAIER

Institut für Physikalische Chemie, Universität Basel, Klingelbergstrasse 80, CH–4056 Basel,
Switzerland,

Neutral unsaturated carbon chains and their ions have been the subject of much recent interest owing to their potential role in the chemistry of the interstellar medium. The detection of spectra for such species and their interpretation is greatly aided by simultaneous theoretical studies.

This poster presents a six–dimensional *ab initio* potential energy surface (PES) for the ground singlet state of C_3H^- generated using the CCSD(T) method and the aug–cc–pVQZ basis set. In agreement with previous studies, the global potential minimum corresponds to a cyclic isomer (C_{2v} , $R_{CC}=1.380$ Å, $R_{CH}=1.083$ Å, $\angle CCH = 46.9^\circ$) and a local minimum is also located, about 2500 cm^{-1} higher in energy, corresponding to a planar trans bent isomer (C_s , $R_{CC}=1.363$ Å, $R_{CH}=1.110$ Å, $\angle CCH = 171.7^\circ$). The PES, represented by a fitted analytical form in the region of the two minima, is used in variational calculations to determine anharmonic rovibrational term values for both isomers and their deuterated analogues. Rotational, centrifugal distortion and vibration–rotation constants are also evaluated.